

What is Addiction? ASAM American Association of Medicine definition of Addiction is a primary, chronic disease of brain reward, motivation, memory and related circuitry. Dysfunction in these circuits leads to characteristic biological, psychological, social and spiritual manifestations. This is reflected in an individual pathologically pursuing reward and/or relief by substance use and other behaviors.


Recovery from addiction is best achieved through a combination of self-management, mutual support, and professional care provided by trained and certified professionals.

Addiction is characterized by inability to consistently abstain, impairment in behavioral control, and craving, diminished recognition of significant problems with one's behaviors and interpersonal relationships, and a dysfunctional emotional response. Like other chronic diseases, addiction often involves cycles of relapse and remission. Without treatment or engagement in recovery activities, addiction is progressive and can result in disability or premature death.

Long Definition of Addiction:

Addiction is a primary, chronic disease of brain reward, motivation, memory and related circuitry. Addiction affects neurotransmission and interactions within reward structures of the brain, including the nucleus accumbens, anterior cingulate cortex, basal forebrain and amygdala, such that motivational hierarchies are altered and addictive behaviors, which may or may not include alcohol and other drug use, supplant healthy, self-care related behaviors. Addiction also affects neurotransmission and interactions between cortical and hippocampal circuits and brain reward structures, such that the memory of previous exposures to rewards (such as food, sex, alcohol and other drugs) leads to a biological and behavioral response to external cues, in turn triggering craving and/or engagement in addictive behaviors.

The neurobiology of addiction encompasses more than the neurochemistry of reward.



The frontal cortex of the brain and underlying white matter connections between the frontal cortex and circuits of reward, motivation and memory are fundamental in the manifestations of altered impulse control, altered judgment, and the dysfunctional pursuit of rewards (which is often experienced by the affected person as a desire to “be normal”) seen in addiction--despite cumulative adverse consequences experienced from engagement in substance use and other addictive behaviors. The frontal lobes are important in inhibiting impulsivity and in assisting individuals to appropriately delay gratification. When persons with addiction manifest problems in deferring gratification, there is a neurological locus of these problems in the frontal cortex. Frontal lobe morphology, connectivity and functioning are still in the process of maturation during adolescence and young adulthood, and early exposure to substance use is another significant factor in the development of addiction. Many neuroscientists

believe that developmental morphology is the basis that makes early-life exposure to substances such an important factor.

Genetic factors account for about half of the likelihood that an individual will develop addiction. Environmental factors interact with the person's biology and affect the extent to which genetic factors exert their influence. Resiliencies the individual acquires (through parenting or later life experiences) can affect the extent to which genetic predispositions lead to the behavioral and other manifestations of addiction. Culture also plays a role in how addiction becomes actualized in persons with biological vulnerabilities to the development of addiction.

Other factors that can contribute to the appearance of addiction, leading to its characteristic bio-psycho-socio-spiritual manifestations, include:

- a. The presence of an underlying biological deficit in the function of reward circuits, such that drugs and behaviors which enhance reward function are preferred and sought as reinforcers;
- b. The repeated engagement in drug use or other addictive behaviors, causing neuroadaptation in motivational circuitry leading to impaired control over further drug use or engagement in addictive behaviors;
- c. Cognitive and affective distortions, which impair perceptions and compromise the ability to deal with feelings, resulting in significant self-deception;
- d. Disruption of healthy social supports and problems in interpersonal relationships which impact the development or impact of resiliencies;
- e. Exposure to trauma or stressors that overwhelm an individual's coping abilities;
- f. Distortion in meaning, purpose and values that guide attitudes, thinking and behavior;
- g. Distortions in a person's connection with self, with others and with the transcendent (referred to as God by many, the Higher Power by 12-steps groups, or higher consciousness by others); and
- h. The presence of co-occurring psychiatric disorders in persons who engage in substance use or other addictive behaviors.

Addiction is characterized by²:

- a. **Inability to consistently Abstain;**
- b. **Impairment in Behavioral control;**
- c. **Craving; or increased “hunger” for drugs or rewarding experiences;**
- d. **Diminished recognition of significant problems with one’s behaviors and interpersonal relationships; and**
- e. **A dysfunctional Emotional response.**

The **power of external cues** to trigger craving and drug use, as well as to increase the frequency of engagement in other potentially addictive behaviors, is also a characteristic of addiction, with the hippocampus being important in memory of previous euphoric or dysphoric experiences, and with the amygdala being important in having motivation concentrate on selecting behaviors associated with these past experiences.

Although some believe that the difference between those who have addiction, and those who do not, is the *quantity* or *frequency* of alcohol/drug use, engagement in addictive behaviors (such as gambling or spending)³, or exposure to other external rewards (such as food or sex), a characteristic aspect of addiction is the *qualitative way* in which the individual responds to such exposures, stressors and environmental cues. A particularly pathological aspect of *the way* that persons with addiction pursue substance use or external rewards is that preoccupation with, obsession with and/or pursuit of rewards (e.g., alcohol and other drug use) persist despite the accumulation of adverse consequences. These manifestations can occur compulsively or impulsively, as a reflection of impaired control.

Persistent risk and/or recurrence of relapse, after periods of abstinence, is another fundamental feature of addiction. This can be triggered by exposure to rewarding substances and behaviors, by exposure to environmental cues to use, and by exposure to emotional stressors that trigger heightened activity in brain stress circuits.⁴

In addiction there is a significant impairment in executive functioning, which manifests in problems with perception, learning, impulse control, compulsivity, and judgment. People with addiction often manifest a lower readiness to change their dysfunctional behaviors despite mounting concerns expressed by significant others in their lives; and display an apparent lack of appreciation of the magnitude of cumulative problems and complications. The still developing frontal

lobes of adolescents may both compound these deficits in executive functioning and predispose youngsters to engage in “high risk” behaviors, including engaging in alcohol or other drug use. The profound drive or craving to use substances or engage in apparently rewarding behaviors, which is seen in many patients with addiction, underscores the compulsive or avolitional aspect of this disease. This is the connection with “powerlessness” over addiction and “unmanageability” of life, as is described in Step 1 of 12 Steps programs.

Addiction is more than a behavioral disorder. Features of addiction include aspects of a person’s behaviors, cognitions, emotions, and interactions with others, including a person’s ability to relate to members of their family, to members of their community, to their own psychological state, and to things that transcend their daily experience.

Behavioral manifestations and complications of addiction, primarily due to impaired control, can include:

- a. Excessive use and/or engagement in addictive behaviors, at higher frequencies and/or quantities than the person intended, often associated with a persistent desire for and unsuccessful attempts at behavioral control;
- b. Excessive time lost in substance use or recovering from the effects of substance use and/or engagement in addictive behaviors, with significant adverse impact on social and occupational functioning (e.g. the development of interpersonal relationship problems or the neglect of responsibilities at home, school or work);
- c. Continued use and/or engagement in addictive behaviors, despite the presence of persistent or recurrent physical or psychological problems which may have been caused or exacerbated by substance use and/or related addictive behaviors;
- d. A narrowing of the behavioral repertoire focusing on rewards that are part of addiction; and
- e. An apparent lack of ability and/or readiness to take consistent, ameliorative action despite recognition of problems.

Cognitive changes in addiction can include:

- a. Preoccupation with substance use;

- b. Altered evaluations of the relative benefits and detriments associated with drugs or rewarding behaviors; and
- c. The inaccurate belief that problems experienced in one's life are attributable to other causes rather than being a predictable consequence of addiction.

Emotional changes in addiction can include:

- a. Increased anxiety, dysphoria and emotional pain;
- b. Increased sensitivity to stressors associated with the recruitment of brain stress systems, such that "things seem more stressful" as a result; and
- c. Difficulty in identifying feelings, distinguishing between feelings and the bodily sensations of emotional arousal, and describing feelings to other people (sometimes referred to as alexithymia).

The emotional aspects of addiction are quite complex. Some persons use alcohol or other drugs or pathologically pursue other rewards because they are seeking "positive reinforcement" or the creation of a positive emotional state ("euphoria"). Others pursue substance use or other rewards because they have experienced relief from negative emotional states ("dysphoria"), which constitutes "negative reinforcement..